

NICHQ Vanderbilt Assessment Scales

Used for diagnosing ADHD



NICHQ Vanderbilt Assessment Scale—PARENT Informant

Today's Date: _____ Child's Name: _____ Date of Birth: _____

Parent's Name: _____ Parent's Phone Number: _____

Directions: Each rating should be considered in the context of what is appropriate for the age of your child.
When completing this form, please think about your child's behaviors in the past 6 months.

Is this evaluation based on a time when the child was on medication was not on medication not sure?

Symptoms	Never	Occasionally	Often	Very Often
1. Does not pay attention to details or makes careless mistakes with, for example, homework	0	1	2	3
2. Has difficulty keeping attention to what needs to be done	0	1	2	3
3. Does not seem to listen when spoken to directly	0	1	2	3
4. Does not follow through when given directions and fails to finish activities (not due to refusal or failure to understand)	0	1	2	3
5. Has difficulty organizing tasks and activities	0	1	2	3
6. Avoids, dislikes, or does not want to start tasks that require ongoing mental effort	0	1	2	3
7. Loses things necessary for tasks or activities (toys, assignments, pencils, or books)	0	1	2	3
8. Is easily distracted by noises or other stimuli	0	1	2	3
9. Is forgetful in daily activities	0	1	2	3
10. Fidgets with hands or feet or squirms in seat	0	1	2	3
11. Leaves seat when remaining seated is expected	0	1	2	3
12. Runs about or climbs too much when remaining seated is expected	0	1	2	3
13. Has difficulty playing or beginning quiet play activities	0	1	2	3
14. Is "on the go" or often acts as if "driven by a motor"	0	1	2	3
15. Talks too much	0	1	2	3
16. Blurts out answers before questions have been completed	0	1	2	3
17. Has difficulty waiting his or her turn	0	1	2	3
18. Interrupts or intrudes in on others' conversations and/or activities	0	1	2	3
19. Argues with adults	0	1	2	3
20. Loses temper	0	1	2	3
21. Actively defies or refuses to go along with adults' requests or rules	0	1	2	3
22. Deliberately annoys people	0	1	2	3
23. Blames others for his or her mistakes or misbehaviors	0	1	2	3
24. Is touchy or easily annoyed by others	0	1	2	3
25. Is angry or resentful	0	1	2	3
26. Is spiteful and wants to get even	0	1	2	3
27. Bullies, threatens, or intimidates others	0	1	2	3
28. Starts physical fights	0	1	2	3
29. Lies to get out of trouble or to avoid obligations (ie, "cons" others)	0	1	2	3
30. Is truant from school (skips school) without permission	0	1	2	3
31. Is physically cruel to people	0	1	2	3
32. Has stolen things that have value	0	1	2	3

The information contained in this publication should not be used as a substitute for the medical care and advice of your pediatrician. There may be variations in treatment that your pediatrician may recommend based on individual facts and circumstances.

Copyright ©2002 American Academy of Pediatrics and National Initiative for Children's Healthcare Quality

Adapted from the Vanderbilt Rating Scales developed by Mark L. Wolraich, MD.

Revised - 1102

American Academy
of Pediatrics



DEDICATED TO THE HEALTH OF ALL CHILDREN™

NICHQ
National Institute for
Children's Health Quality



NICHQ Vanderbilt Assessment Scale—PARENT Informant

Today's Date: _____ Child's Name: _____ Date of Birth: _____

Parent's Name: _____ Parent's Phone Number: _____

Symptoms (continued)	Never	Occasionally	Often	Very Often
33. Deliberately destroys others' property	0	1	2	3
34. Has used a weapon that can cause serious harm (bat, knife, brick, gun)	0	1	2	3
35. Is physically cruel to animals	0	1	2	3
36. Has deliberately set fires to cause damage	0	1	2	3
37. Has broken into someone else's home, business, or car	0	1	2	3
38. Has stayed out at night without permission	0	1	2	3
39. Has run away from home overnight	0	1	2	3
40. Has forced someone into sexual activity	0	1	2	3
41. Is fearful, anxious, or worried	0	1	2	3
42. Is afraid to try new things for fear of making mistakes	0	1	2	3
43. Feels worthless or inferior	0	1	2	3
44. Blames self for problems, feels guilty	0	1	2	3
45. Feels lonely, unwanted, or unloved; complains that "no one loves him or her"	0	1	2	3
46. Is sad, unhappy, or depressed	0	1	2	3
47. Is self-conscious or easily embarrassed	0	1	2	3

Performance	Excellent	Above Average	Average	Somewhat of a Problem	Problematic
48. Overall school performance	1	2	3	4	5
49. Reading	1	2	3	4	5
50. Writing	1	2	3	4	5
51. Mathematics	1	2	3	4	5
52. Relationship with parents	1	2	3	4	5
53. Relationship with siblings	1	2	3	4	5
54. Relationship with peers	1	2	3	4	5
55. Participation in organized activities (eg, teams)	1	2	3	4	5

Comments:

For Office Use Only

Total number of questions scored 2 or 3 in questions 1–9: _____

Total number of questions scored 2 or 3 in questions 10–18: _____

Total Symptom Score for questions 1–18: _____

Total number of questions scored 2 or 3 in questions 19–26: _____

Total number of questions scored 2 or 3 in questions 27–40: _____

Total number of questions scored 2 or 3 in questions 41–47: _____

Total number of questions scored 4 or 5 in questions 48–55: _____

Average Performance Score: _____



Teacher's Name: _____ Class Time: _____ Class Name/Period: _____

Today's Date: _____ Child's Name: _____ Grade Level: _____

Directions: Each rating should be considered in the context of what is appropriate for the age of the child you are rating and should reflect that child's behavior since the beginning of the school year. Please indicate the number of weeks or months you have been able to evaluate the behaviors: _____.

Is this evaluation based on a time when the child was on medication was not on medication not sure?

Symptoms	Never	Occasionally	Often	Very Often
1. Fails to give attention to details or makes careless mistakes in schoolwork	0	1	2	3
2. Has difficulty sustaining attention to tasks or activities	0	1	2	3
3. Does not seem to listen when spoken to directly	0	1	2	3
4. Does not follow through on instructions and fails to finish schoolwork (not due to oppositional behavior or failure to understand)	0	1	2	3
5. Has difficulty organizing tasks and activities	0	1	2	3
6. Avoids, dislikes, or is reluctant to engage in tasks that require sustained mental effort	0	1	2	3
7. Loses things necessary for tasks or activities (school assignments, pencils, or books)	0	1	2	3
8. Is easily distracted by extraneous stimuli	0	1	2	3
9. Is forgetful in daily activities	0	1	2	3
10. Fidgets with hands or feet or squirms in seat	0	1	2	3
11. Leaves seat in classroom or in other situations in which remaining seated is expected	0	1	2	3
12. Runs about or climbs excessively in situations in which remaining seated is expected	0	1	2	3
13. Has difficulty playing or engaging in leisure activities quietly	0	1	2	3
14. Is "on the go" or often acts as if "driven by a motor"	0	1	2	3
15. Talks excessively	0	1	2	3
16. Blurts out answers before questions have been completed	0	1	2	3
17. Has difficulty waiting in line	0	1	2	3
18. Interrupts or intrudes on others (eg, butts into conversations/games)	0	1	2	3
19. Loses temper	0	1	2	3
20. Actively defies or refuses to comply with adult's requests or rules	0	1	2	3
21. Is angry or resentful	0	1	2	3
22. Is spiteful and vindictive	0	1	2	3
23. Bullies, threatens, or intimidates others	0	1	2	3
24. Initiates physical fights	0	1	2	3
25. Lies to obtain goods for favors or to avoid obligations (eg, "cons" others)	0	1	2	3
26. Is physically cruel to people	0	1	2	3
27. Has stolen items of nontrivial value	0	1	2	3
28. Deliberately destroys others' property	0	1	2	3
29. Is fearful, anxious, or worried	0	1	2	3
30. Is self-conscious or easily embarrassed	0	1	2	3
31. Is afraid to try new things for fear of making mistakes	0	1	2	3

The recommendations in this publication do not indicate an exclusive course of treatment or serve as a standard of medical care. Variations, taking into account individual circumstances, may be appropriate.

Copyright ©2002 American Academy of Pediatrics and National Initiative for Children's Healthcare Quality

Adapted from the Vanderbilt Rating Scales developed by Mark L. Wolraich, MD.

Revised - 0303

American Academy
of Pediatrics



DEDICATED TO THE HEALTH OF ALL CHILDREN™

NICHQ
National Institute for
Children's Health Quality



HE0351

Teacher's Name: _____ Class Time: _____ Class Name/Period: _____

Today's Date: _____ Child's Name: _____ Grade Level: _____

Symptoms (continued)	Never	Occasionally	Often	Very Often
32. Feels worthless or inferior	0	1	2	3
33. Blames self for problems; feels guilty	0	1	2	3
34. Feels lonely, unwanted, or unloved; complains that "no one loves him or her"	0	1	2	3
35. Is sad, unhappy, or depressed	0	1	2	3

Performance	Excellent	Above Average	Average	Somewhat of a Problem	Problematic
Academic Performance					
36. Reading	1	2	3	4	5
37. Mathematics	1	2	3	4	5
38. Written expression	1	2	3	4	5

Classroom Behavioral Performance	Excellent	Above Average	Average	Somewhat of a Problem	Problematic
39. Relationship with peers	1	2	3	4	5
40. Following directions	1	2	3	4	5
41. Disrupting class	1	2	3	4	5
42. Assignment completion	1	2	3	4	5
43. Organizational skills	1	2	3	4	5

Comments:

Please return this form to: _____

Mailing address: _____

Fax number: _____

For Office Use Only

Total number of questions scored 2 or 3 in questions 1–9: _____

Total number of questions scored 2 or 3 in questions 10–18: _____

Total Symptom Score for questions 1–18: _____

Total number of questions scored 2 or 3 in questions 19–28: _____

Total number of questions scored 2 or 3 in questions 29–35: _____

Total number of questions scored 4 or 5 in questions 36–43: _____

Average Performance Score: _____

American Academy
of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™

11-20/rev0303

NICHQ
National Institute for
Children's Health Quality

Today's Date: _____ Child's Name: _____ Date of Birth: _____

Parent's Name: _____ Parent's Phone Number: _____

Directions: Each rating should be considered in the context of what is appropriate for the age of your child. Please think about your child's behaviors since the last assessment scale was filled out when rating his/her behaviors.

Is this evaluation based on a time when the child was on medication was not on medication not sure?

Symptoms	Never	Occasionally	Often	Very Often
1. Does not pay attention to details or makes careless mistakes with, for example, homework	0	1	2	3
2. Has difficulty keeping attention to what needs to be done	0	1	2	3
3. Does not seem to listen when spoken to directly	0	1	2	3
4. Does not follow through when given directions and fails to finish activities (not due to refusal or failure to understand)	0	1	2	3
5. Has difficulty organizing tasks and activities	0	1	2	3
6. Avoids, dislikes, or does not want to start tasks that require ongoing mental effort	0	1	2	3
7. Loses things necessary for tasks or activities (toys, assignments, pencils, or books)	0	1	2	3
8. Is easily distracted by noises or other stimuli	0	1	2	3
9. Is forgetful in daily activities	0	1	2	3
10. Fidgets with hands or feet or squirms in seat	0	1	2	3
11. Leaves seat when remaining seated is expected	0	1	2	3
12. Runs about or climbs too much when remaining seated is expected	0	1	2	3
13. Has difficulty playing or beginning quiet play activities	0	1	2	3
14. Is "on the go" or often acts as if "driven by a motor"	0	1	2	3
15. Talks too much	0	1	2	3
16. Blurts out answers before questions have been completed	0	1	2	3
17. Has difficulty waiting his or her turn	0	1	2	3
18. Interrupts or intrudes in on others' conversations and/or activities	0	1	2	3

Performance	Excellent	Above Average	Average	Somewhat of a Problem	Problematic
19. Overall school performance	1	2	3	4	5
20. Reading	1	2	3	4	5
21. Writing	1	2	3	4	5
22. Mathematics	1	2	3	4	5
23. Relationship with parents	1	2	3	4	5
24. Relationship with siblings	1	2	3	4	5
25. Relationship with peers	1	2	3	4	5
26. Participation in organized activities (eg, teams)	1	2	3	4	5

The information contained in this publication should not be used as a substitute for the medical care and advice of your pediatrician. There may be variations in treatment that your pediatrician may recommend based on individual facts and circumstances.

Copyright ©2002 American Academy of Pediatrics and National Initiative for Children's Healthcare Quality

Adapted from the Vanderbilt Rating Scales developed by Mark L. Wolraich, MD.

Revised - 0303

American Academy
of Pediatrics



DEDICATED TO THE HEALTH OF ALL CHILDREN™

NICHQ
National Institute for
Children's Health Quality



Today's Date: _____ Child's Name: _____ Date of Birth: _____

Parent's Name: _____ Parent's Phone Number: _____

Side Effects: Has your child experienced any of the following side effects or problems in the past week?	Are these side effects currently a problem?			
	None	Mild	Moderate	Severe
Headache				
Stomachache				
Change of appetite—explain below				
Trouble sleeping				
Irritability in the late morning, late afternoon, or evening—explain below				
Socially withdrawn—decreased interaction with others				
Extreme sadness or unusual crying				
Dull, tired, listless behavior				
Tremors/feeling shaky				
Repetitive movements, tics, jerking, twitching, eye blinking—explain below				
Picking at skin or fingers, nail biting, lip or cheek chewing—explain below				
Sees or hears things that aren't there				

Explain/Comments:

For Office Use Only

Total Symptom Score for questions 1–18: _____

Average Performance Score for questions 19–26: _____

Adapted from the Pittsburgh side effects scale, developed by William E. Pelham, Jr, PhD.

American Academy
of Pediatrics



DEDICATED TO THE HEALTH OF ALL CHILDREN™

NICHQ
National Institute for
Children's Health Quality

McNeil
Consumer & Specialty Pharmaceuticals

Teacher's Name: _____ Class Time: _____ Class Name/Period: _____

Today's Date: _____ Child's Name: _____ Grade Level: _____

Directions: Each rating should be considered in the context of what is appropriate for the age of the child you are rating and should reflect that child's behavior since the last assessment scale was filled out. Please indicate the number of weeks or months you have been able to evaluate the behaviors: _____.

Is this evaluation based on a time when the child was on medication was not on medication not sure?

Symptoms	Never	Occasionally	Often	Very Often
1. Does not pay attention to details or makes careless mistakes with, for example, homework	0	1	2	3
2. Has difficulty keeping attention to what needs to be done	0	1	2	3
3. Does not seem to listen when spoken to directly	0	1	2	3
4. Does not follow through when given directions and fails to finish activities (not due to refusal or failure to understand)	0	1	2	3
5. Has difficulty organizing tasks and activities	0	1	2	3
6. Avoids, dislikes, or does not want to start tasks that require ongoing mental effort	0	1	2	3
7. Loses things necessary for tasks or activities (toys, assignments, pencils, or books)	0	1	2	3
8. Is easily distracted by noises or other stimuli	0	1	2	3
9. Is forgetful in daily activities	0	1	2	3
10. Fidgets with hands or feet or squirms in seat	0	1	2	3
11. Leaves seat when remaining seated is expected	0	1	2	3
12. Runs about or climbs too much when remaining seated is expected	0	1	2	3
13. Has difficulty playing or beginning quiet play activities	0	1	2	3
14. Is "on the go" or often acts as if "driven by a motor"	0	1	2	3
15. Talks too much	0	1	2	3
16. Blurts out answers before questions have been completed	0	1	2	3
17. Has difficulty waiting his or her turn	0	1	2	3
18. Interrupts or intrudes in on others' conversations and/or activities	0	1	2	3

Performance	Somewhat of a Problem				
	Excellent	Above Average	Average	Problem	Problematic
19. Reading	1	2	3	4	5
20. Mathematics	1	2	3	4	5
21. Written expression	1	2	3	4	5
22. Relationship with peers	1	2	3	4	5
23. Following direction	1	2	3	4	5
24. Disrupting class	1	2	3	4	5
25. Assignment completion	1	2	3	4	5
26. Organizational skills	1	2	3	4	5

The recommendations in this publication do not indicate an exclusive course of treatment or serve as a standard of medical care. Variations, taking into account individual circumstances, may be appropriate.

Copyright ©2002 American Academy of Pediatrics and National Initiative for Children's Healthcare Quality

Adapted from the Vanderbilt Rating Scales developed by Mark L. Wolraich, MD.

Revised - 0303

American Academy
of Pediatrics



DEDICATED TO THE HEALTH OF ALL CHILDREN™

NICHQ
National Institute for
Children's Health Quality

McNeil
Consumer & Specialty Pharmaceuticals

Teacher's Name: _____ Class Time: _____ Class Name/Period: _____

Today's Date: _____ Child's Name: _____ Grade Level: _____

Side Effects: Has the child experienced any of the following side effects or problems in the past week?	Are these side effects currently a problem?			
	None	Mild	Moderate	Severe
Headache				
Stomachache				
Change of appetite—explain below				
Trouble sleeping				
Irritability in the late morning, late afternoon, or evening—explain below				
Socially withdrawn—decreased interaction with others				
Extreme sadness or unusual crying				
Dull, tired, listless behavior				
Tremors/feeling shaky				
Repetitive movements, tics, jerking, twitching, eye blinking—explain below				
Picking at skin or fingers, nail biting, lip or cheek chewing—explain below				
Sees or hears things that aren't there				

Explain/Comments:**For Office Use Only**

Total Symptom Score for questions 1–18: _____

Average Performance Score: _____

Please return this form to: _____

Mailing address: _____

Fax number: _____

Adapted from the Pittsburgh side effects scale, developed by William E. Pelham, Jr, PhD.

American Academy
of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™

11-22/rev0303

NICHQ
National Institute for
Children's Health QualityMcNeil
Consumer & Specialty Pharmaceuticals

Scoring Instructions for the NICHQ Vanderbilt Assessment Scales

These scales should NOT be used alone to make any diagnosis. You must take into consideration information from multiple sources. Scores of 2 or 3 on a single Symptom question reflect **often-occurring** behaviors. Scores of 4 or 5 on Performance questions reflect problems in performance.

The initial assessment scales, parent and teacher, have 2 components: symptom assessment and impairment in performance. On both the parent and teacher initial scales, the symptom assessment screens for symptoms that meet criteria for both inattentive (items 1–9) and hyperactive ADHD (items 10–18).

To meet *DSM-IV* criteria for the diagnosis, one must have at least 6 positive responses to either the inattentive 9 or hyperactive 9 core symptoms, or both. A positive response is a 2 or 3 (often, very often) (you could draw a line straight down the page and count the positive answers in each subsegment). There is a place to

record the number of positives in each subsegment, and a place for total score for the first 18 symptoms (just add them up).

The initial scales also have symptom screens for 3 other co-morbidities—oppositional-defiant, conduct, and anxiety/depression. These are screened by the number of positive responses in each of the segments separated by the “squares.” The specific item sets and numbers of positives required for each co-morbid symptom screen set are detailed below.

The second section of the scale has a set of performance measures, scored 1 to 5, with 4 and 5 being somewhat of a problem/problematic. To meet criteria for ADHD there must be at least one item of the Performance set in which the child scores a 4 or 5; ie, there must be impairment, not just symptoms to meet diagnostic criteria. The sheet has a place to record the number of positives (4s, 5s) and an Average Performance Score—add them up and divide by number of Performance criteria answered.

Parent Assessment Scale	Teacher Assessment Scale
<p>Predominantly Inattentive subtype</p> <ul style="list-style-type: none"> ■ Must score a 2 or 3 on 6 out of 9 items on questions 1–9 <u>AND</u> ■ Score a 4 or 5 on any of the Performance questions 48–55 <p>Predominantly Hyperactive/Impulsive subtype</p> <ul style="list-style-type: none"> ■ Must score a 2 or 3 on 6 out of 9 items on questions 10–18 <u>AND</u> ■ Score a 4 or 5 on any of the Performance questions 48–55 <p>ADHD Combined Inattention/Hyperactivity</p> <ul style="list-style-type: none"> ■ Requires the above criteria on both inattention and hyperactivity/impulsivity <p>Oppositional-Defiant Disorder Screen</p> <ul style="list-style-type: none"> ■ Must score a 2 or 3 on 4 out of 8 behaviors on questions 19–26 <u>AND</u> ■ Score a 4 or 5 on any of the Performance questions 48–55 <p>Conduct Disorder Screen</p> <ul style="list-style-type: none"> ■ Must score a 2 or 3 on 3 out of 14 behaviors on questions 27–40 <u>AND</u> ■ Score a 4 or 5 on any of the Performance questions 48–55 <p>Anxiety/Depression Screen</p> <ul style="list-style-type: none"> ■ Must score a 2 or 3 on 3 out of 7 behaviors on questions 41–47 <u>AND</u> ■ Score a 4 or 5 on any of the Performance questions 48–55 	<p>Predominantly Inattentive subtype</p> <ul style="list-style-type: none"> ■ Must score a 2 or 3 on 6 out of 9 items on questions 1–9 <u>AND</u> ■ Score a 4 or 5 on any of the Performance questions 36–43 <p>Predominantly Hyperactive/Impulsive subtype</p> <ul style="list-style-type: none"> ■ Must score a 2 or 3 on 6 out of 9 items on questions 10–18 <u>AND</u> ■ Score a 4 or 5 on any of the Performance questions 36–43 <p>ADHD Combined Inattention/Hyperactivity</p> <ul style="list-style-type: none"> ■ Requires the above criteria on both inattention and hyperactivity/impulsivity <p>Oppositional-Defiant/Conduct Disorder Screen</p> <ul style="list-style-type: none"> ■ Must score a 2 or 3 on 3 out of 10 items on questions 19–28 <u>AND</u> ■ Score a 4 or 5 on any of the Performance questions 36–43 <p>Anxiety/Depression Screen</p> <ul style="list-style-type: none"> ■ Must score a 2 or 3 on 3 out of 7 items on questions 29–35 <u>AND</u> ■ Score a 4 or 5 on any of the Performance questions 36–43

The parent and teacher follow-up scales have the first 18 core ADHD symptoms, not the co-morbid symptoms. The section segment has the same Performance items and impairment assessment as the initial scales, and then has a side-effect reporting scale that can be used to both assess and monitor the presence of adverse reactions to medications prescribed, if any.

Scoring the follow-up scales involves only calculating a total symptom score for items 1–18 that can be tracked over time, and

the average of the Performance items answered as measures of improvement over time with treatment.

Parent Assessment Follow-up

- Calculate Total Symptom Score for questions 1–18.
- Calculate Average Performance Score for questions 19–26.

Teacher Assessment Follow-up

- Calculate Total Symptom Score for questions 1–18.
- Calculate Average Performance Score for questions 19–26.

The recommendations in this publication do not indicate an exclusive course of treatment or serve as a standard of medical care. Variations, taking into account individual circumstances, may be appropriate.

Copyright ©2002 American Academy of Pediatrics and National Initiative for Children's Healthcare Quality

American Academy
of Pediatrics



DEDICATED TO THE HEALTH OF ALL CHILDREN™

NICHQ
National Institute for
Children's Health Quality



Adult ADHD Self-Report Scale (ASRS-v1.1) Symptom Checklist Instructions

The questions on the back page are designed to stimulate dialogue between you and your patients and to help confirm if they may be suffering from the symptoms of attention-deficit/hyperactivity disorder (ADHD).

Description: The Symptom Checklist is an instrument consisting of the eighteen DSM-IV-TR criteria. Six of the eighteen questions were found to be the most predictive of symptoms consistent with ADHD. These six questions are the basis for the ASRS v1.1 Screener and are also Part A of the Symptom Checklist. Part B of the Symptom Checklist contains the remaining twelve questions.

Instructions:

Symptoms

1. Ask the patient to complete both Part A and Part B of the Symptom Checklist by marking an X in the box that most closely represents the frequency of occurrence of each of the symptoms.
2. Score Part A. If four or more marks appear in the darkly shaded boxes within Part A then the patient has symptoms highly consistent with ADHD in adults and further investigation is warranted.
3. The frequency scores on Part B provide additional cues and can serve as further probes into the patient's symptoms. Pay particular attention to marks appearing in the dark shaded boxes. The frequency-based response is more sensitive with certain questions. No total score or diagnostic likelihood is utilized for the twelve questions. It has been found that the six questions in Part A are the most predictive of the disorder and are best for use as a screening instrument.

Impairments

1. Review the entire Symptom Checklist with your patients and evaluate the level of impairment associated with the symptom.
2. Consider work/school, social and family settings.
3. Symptom frequency is often associated with symptom severity, therefore the Symptom Checklist may also aid in the assessment of impairments. If your patients have frequent symptoms, you may want to ask them to describe how these problems have affected the ability to work, take care of things at home, or get along with other people such as their spouse/significant other.

History

1. Assess the presence of these symptoms or similar symptoms in childhood. Adults who have ADHD need not have been formally diagnosed in childhood. In evaluating a patient's history, look for evidence of early-appearing and long-standing problems with attention or self-control. Some significant symptoms should have been present in childhood, but full symptomology is not necessary.

Adult ADHD Self-Report Scale (ASRS-v1.1) Symptom Checklist

Patient Name		Today's Date					
Please answer the questions below, rating yourself on each of the criteria shown using the scale on the right side of the page. As you answer each question, place an X in the box that best describes how you have felt and conducted yourself over the past 6 months. Please give this completed checklist to your healthcare professional to discuss during today's appointment.			Never	Rarely	Sometimes	Often	Very Often
1. How often do you have trouble wrapping up the final details of a project, once the challenging parts have been done?							
2. How often do you have difficulty getting things in order when you have to do a task that requires organization?							
3. How often do you have problems remembering appointments or obligations?							
4. When you have a task that requires a lot of thought, how often do you avoid or delay getting started?							
5. How often do you fidget or squirm with your hands or feet when you have to sit down for a long time?							
6. How often do you feel overly active and compelled to do things, like you were driven by a motor?							
Part A							
7. How often do you make careless mistakes when you have to work on a boring or difficult project?							
8. How often do you have difficulty keeping your attention when you are doing boring or repetitive work?							
9. How often do you have difficulty concentrating on what people say to you, even when they are speaking to you directly?							
10. How often do you misplace or have difficulty finding things at home or at work?							
11. How often are you distracted by activity or noise around you?							
12. How often do you leave your seat in meetings or other situations in which you are expected to remain seated?							
13. How often do you feel restless or fidgety?							
14. How often do you have difficulty unwinding and relaxing when you have time to yourself?							
15. How often do you find yourself talking too much when you are in social situations?							
16. When you're in a conversation, how often do you find yourself finishing the sentences of the people you are talking to, before they can finish them themselves?							
17. How often do you have difficulty waiting your turn in situations when turn taking is required?							
18. How often do you interrupt others when they are busy?							
Part B							

The Value of Screening for Adults With ADHD

Research suggests that the symptoms of ADHD can persist into adulthood, having a significant impact on the relationships, careers, and even the personal safety of your patients who may suffer from it.¹⁻⁴ Because this disorder is often misunderstood, many people who have it do not receive appropriate treatment and, as a result, may never reach their full potential. Part of the problem is that it can be difficult to diagnose, particularly in adults.

The Adult ADHD Self-Report Scale (ASRS-v1.1) Symptom Checklist was developed in conjunction with the World Health Organization (WHO), and the Workgroup on Adult ADHD that included the following team of psychiatrists and researchers:

- **Lenard Adler, MD**
Associate Professor of Psychiatry and Neurology
New York University Medical School
- **Ronald C. Kessler, PhD**
Professor, Department of Health Care Policy
Harvard Medical School
- **Thomas Spencer, MD**
Associate Professor of Psychiatry
Harvard Medical School

As a healthcare professional, you can use the ASRS v1.1 as a tool to help screen for ADHD in adult patients. Insights gained through this screening may suggest the need for a more in-depth clinician interview. The questions in the ASRS v1.1 are consistent with DSM-IV criteria and address the manifestations of ADHD symptoms in adults. Content of the questionnaire also reflects the importance that DSM-IV places on symptoms, impairments, and history for a correct diagnosis.⁴

The checklist takes about 5 minutes to complete and can provide information that is critical to supplement the diagnostic process.

References:

1. Schweitzer JB, et al. *Med Clin North Am.* 2001;85(3):10-11, 757-777.
2. Barkley RA. *Attention Deficit Hyperactivity Disorder: A Handbook for Diagnosis and Treatment.* 2nd ed. 1998.
3. Biederman J, et al. *Am J Psychiatry.* 1993;150:1792-1798.
4. American Psychiatric Association: *Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition, Text Revision.* Washington, DC, American Psychiatric Association. 2000: 85-93.